

CRYPTOBACTERIUM CURTUM: A NOVEL BACTERIA WITH A POSSIBLE SIGNIFICANT INVOLVEMENT IN PERIODONTAL DISEASE. A REVIEW.

Oana Bejan¹, Giorgio Nichitean^{2*}, Oana Butnaru^{3*}, Monica Tatarciuc⁴, Maria-Alexandra Martu⁵, Gabriel Rotundu⁵, Iris-Madalina Maftai⁶, Costin Damian⁷, Ramona Ursu⁷, Ionuț Luchian⁵

¹“Gr.T.Popa”, U.M.Ph. - Iași, Romania, Faculty of Dental Medicine, Resident Doctor

² PhD Student “Gr.T.Popa” U.M.Ph. - Iași, Romania

³“Gr.T.Popa”, U.M.Ph. - Iași, Romania, Faculty of Dental Medicine, Department of Biophysics

⁴“Gr.T.Popa”, U.M.Ph. - Iași, Romania, Faculty of Dental Medicine, Department of Dental Technology

⁵“Gr.T.Popa”, U.M.Ph. - Iași, Romania, Faculty of Dental Medicine, Department of Periodontology

⁶“Gr.T.Popa”, U.M.Ph. - Iași, Romania, Faculty of Dental Medicine. Student

⁷“Gr.T.Popa”, U.M.Ph. - Iași, Romania, Faculty of Medicine, Department of Microbiology

*Corresponding authors: Giorgio Nichitean: nichitean.marian-giorgio@d.umfiasi.ro

Oana Butnaru: oana.maria.butnaru@umfiasi.ro

all authors had an equal contribution as the first authors

ABSTRACT

Periodontal disease is considered to be connected with some very well-known anaerobic etiological agents. Many studies are analysing the associations between Porphyromonas gingivalis, Aggregatibacter actinomycetemcomitans and Tannerella forsythia with periodontal disease. Still, the prevalence of this disease is increased. In this review we aimed to present a possible new bacteria connected with periodontal disease: Cryptobacterium curtum. We performed a search in PubMed and we identified 8 articles, published between 1999 and 2018. Here we present a summary about this bacterial species regarding its microscopic, cultivation and genomic features. New molecular biology techniques, such as NGS (next generation sequencing) or mRNA technology, identified this bacteria to be associated with acute dental abscesses, and with chronic periodontitis. The virulence factors of this bacteria are not yet identified, but with the recent highly sensitive techniques, there is hope to completely establish the role of C. curtum in the pathogenesis of periodontal disease.

Keywords: *periodontal disease, etiological agents, modern technology, NGS, mRNA*

1. INTRODUCTION

Periodontitis is a polymicrobial disease caused by complex interactions between distinct pathogens.[1] At the biofilm level we can observe the occurrence of destructive phenomena on periodontal tissues.[2,3] It is unanimously accepted in the literature that certain microorganisms, which are still poorly understood, may be involved in the onset or progression of periodontitis [4]. For several decades, oral microbiology research has failed to identify

the composition of the subgingival microbiota due to technical limitations.[5]

In the last decade, due to paradigm shifts involving the use of molecular biology and sequencing techniques, the detection of novel periodontal pathogens has become feasible.[4-6] It is therefore evident that, in addition to conventional periodontal pathogens, other microorganisms could be significantly involved in the occurrence and progression of periodontal and peri-implant pathological processes.[7,8]

Among the new periodontopathogenic microbial agents we can list *Cryptobacterium curtum*, *Dialister pneumosintes*, *Filifactor alocis*, *Mitsuokella dentalis*, *Slackia exigua*, *Selenomonas sputigena*, *Solobacterium moorei*, *Treponema lecithinolyticum* and *Synergistes*. [9]

2. CRYPTOBACTERIUM CURTUM - GENERALITIES

The name *Cryptobacterium curtum* derives from the Greek words "Kryptos," which means "hidden" and "curtum," which means "shortened." These aspects led to its inclusion in the genus *Eubacterium*, but subsequent reclassification was necessary into a new genus called *Cryptobacterium*. [10-11] We can point out that it belongs to the domain *Bacteria*, phylum *Actinobacteria*, class *Actinobacteria*, order *Coriobacteriales*, family *Coriobacteriaceae*, genus *Cryptobacterium*, and species *Curtum*. Morphologically, they exhibit characteristics pathognomonic of short gram-positive bacilli, obligate anaerobic, lacking motility and spores, mesophilic, non-sucrolytic and can range in size from 0.8 μm to 1.0 μm . Ultrathin sections showed a single-layered cell wall, with a thickness of 10 nm, in which no pili or flagella were identified (Fig. 1).




Figure 1. Schematic representation of *Cryptobacterium curtum*

On specific and favourable culture media, the formation of small translucent non-haemolytic colonies with a < 1 mm (0.3-0.5 mm) diameter can be observed. [9-11]

Growth of *C. curtum* in broth with glucose-supplemented peptone yeast extract can be optionally augmented with 0.5% arginine, derived by enzymatic degradation of arginine deaminase-associated peptides. At the same time, *C. curtum* shows poor growth due to limited substrate utilisation for metabolic activity. [10-13]

Cryptobacterium curtum is a relatively recently identified and characterized species that has been isolated from the gingival sulcus of patients with periodontal disease. [14] There is, however, limited scientific evidence demonstrating more frequent detection of *T. socranskii* in human subjects with periodontitis, although the observed differences were not statistically significant. [15] At the same time, we can recall that the prevalence of the bacterium was higher with the increasing of probing depth and the exceeding of the physiological threshold. [16]

3. DESCRIPTION OF CRYPTOBACTERIUM GEN. NOV.

Cryptobacterium (Crypt. 0.b ac. te'ri.um. Gr. n. kryptos hiding; Gr. n. bakterion a small bacillus; M.L. neut. n. *Cryptobacterium* a bacteria hiding in the form of a bacillus). The cells may present as short Gram-positive bacilli, occasionally Gram-variable, when in the stationary phase. They are obligately anaerobic, lacking motility and non-spore-forming. Biochemically, they are catalase-negative, asaccharolytic and without volatile end products in glucose-supplemented peptone yeast extract. From an analytical

perspective we can mention that the G + C DNA content is 50-51 mol%. According to the almost complete 16s rDNA sequence, the genus groups with the general actinomycete group of Gram-positive bacteria and the type species is *Cryptobacterium curturn*. [14]

4. DESCRIPTION OF CRYPTOBACTERIUM CURTUM SP. NOV.

Cryptobacterium curtum (cur'tum. L. neut. adj. shortened curtum, a shortened cell of this organism). This description is based on the study of two strains isolated from the oral cavities of patients with periodontal disease. The cells are very short bacilli, Gram-positive, obligately anaerobic, non-motile and non-spore-forming. [11-15] The cells may be individual or may appear as arranged in clusters. Older cultures sometimes stain Gram-negative, due to changes that may occur in the bacterial cell wall. Tiny colonies, less than 1 mm in diameter, form on culture media and are circular, convex and translucent even after prolonged incubation in an anaerobic glove box. Growth in broth media is poor with or without carbohydrates. No hemolysis occurs on blood agar plates with brain heart infusion blood. Cells are inert in most biochemical tests. [23] Starch and esculin are not hydrolysed; nitrate is not reduced.

No liquefaction of gelatin occurs. Indole, urease and catalase tests are also negative. Ammonia is produced from arginine. The strains are non-fermentative and do not utilise adonitol, amygdalin, arabinose, cellobiose, erythritol, aesculin, fructose, galactose, glucose, glycogen, inositol, lactose, maltose, mannitol, mannose, melezitose, melibiose, rhamnose, ribose, salicin, sorbitol, starch, sucrose, trehalose or xylose. [24-26]

No metabolic end product is detected in glucose-supplemented peptone yeast extract medium or glucose-supplemented peptone yeast extract broth. It was isolated from the periodontal pocket of a human adult with periodontal disease. The G + C DNA content is 50-51 mol%. [14]

5. GENOME PROPERTIES

The genome has a length of 1 617 804 bp and comprises a main circular chromosome with 50.9% GC content (Table I). Of the 1422 predicted genes, 1364 were protein-coding genes and 58 were RNA-coding genes. [27-29] A total of 7 pseudogenes were also identified. Among the majority of protein-coding genes (78.5%) were assigned with a putative function, while the rest were noted as hypothetical proteins. [30-31] Genome properties and statistics are summarised in Table I. The distribution of genes into COG functional categories is shown in Table II. [10]

Table I: Genes of *Cryptobacterium curtum* identified by complete genome sequencing (adapted from Mavrommatis et al., 2009)

Category of gene	% of Total
RNA genes	2.37%
rRNA operons	
Protein-coding genes	95.92%
Pseudo genes	0.49%
Genes with function prediction	78.55%
Genes in paralog clusters	5.41 %
Genes assigned to COGs	77.57%
Genes assigned Pfam domains	77.64%
Genes with signal peptides	19.37%
Genes with transmembrane helices	14.46%

Table II. The most representative genes of *Cryptobacterium curtum* (adapted from Mavrommatis et al., 2009)

Description of gene function	%
Translation, ribosomal structure and biogenesis	9.4
RNA processing and modification	0.1
Transcription	6.9
Replication, recombination and repair	5.5
Chromatin structure and dynamics	0.1
Cell cycle control, mitosis and meiosis	1.1
Nuclear structure	0.0
Defense mechanisms	1.5
Signal transduction mechanisms	4.7
Cell wall/membrane biogenesis	5.1
Cell motility	0.1
Cytoskeleton	0.1

Other genes identified related to energy production and conversion, carbohydrate transport and metabolism, amino acid

transport and metabolism, nucleotide transport and metabolism, coenzyme transport and metabolism, lipid transport

and metabolism, inorganic ion transport and metabolism. This full complete genome analysis performed with old and new NGS technologies offered the possibility of properly classification of this bacterial species, from *Eubacterium (Firmicutes)* to *Actinobacteria*, close to the *Coriobacteriaceae*. [10]

6. ASSOCIATION OF *C. CURTUM* WITH PERIODONTAL DISEASE

Kumar et al., 2003, [32] using PCR amplification of the 16S rDNA and the downstream intergenic spacer region (ISR) for bacterial species and phylotypes detection, identified *C. curtum* with a prevalence of 33% in healthy persons and with a prevalence of 64% in periodontitis patients, with a $p = 0.0005$, which suggest its significant association as a possible etiologic agent. Using 16S gene sequenced on the Illumina MiSeq platform, Lopez-Oliva et al., 2018, [33] identified *Cryptobacterium curtum*, from subgingival plaque of individuals with periodontal disease. These two very sensitive molecular biology techniques have the advantage of detecting new bacterial species associated with periodontal disease, but only after larger studies very well conducted, would it be possible to confirm its pathogenicity and its virulence factors associated with periodontal diseases.

REFERENCES:

1. Teodorescu AC, Teslaru S, Solomon SM, Zetu L, Luchian I, Sioustis IA, Martu, MA, Vasiliu B, Martu S. Assessment of Bacterial Associations Involved in Periodontal Disease Using Crevicular Fluid. *Rev. Chim.(Bucharest)*, 2019;70(6):2145-2149
2. Martu MA, Solomon SM, Sufaru IG, Jelihovschi I, Martu S, Rezus E, Surdu AE, Onea RM, Grecu GP, Foia L. Study on the Prevalence of Periodontopathogenic Bacteria in Serum and Subgingival Bacterial Plaque in Patients with Rheumatoid Arthritis. *Rev. Chim. (Bucharest)*. 2017; 68(8):1946-1949

The diversity and complexity observed are likely due to several factors, including the interdependence of bacterial species within commensal communities as well as inter-individual variation in microbial aetiology and host susceptibility.[34,35] Elucidation of the pathogenesis mechanisms of periodontal and peri-implant diseases requires the initiation of future studies focusing on obtaining quantitative information on the proportions of these newly identified species especially at active disease sites.[7]

7. CONCLUSION AND PERSPECTIVES

The microbial aetiology of both chronic and acute periodontal and peri-implant processes is more complex than the information provided in the current literature and a large number of bacterial species, appear to show a possible association with disease. This premise necessitated the expansion of the list to include several uncultured species but recently identified, by ribosomal sequence analysis.

Longitudinal studies may play a key role in understanding the natural history of this chronic disease. At the same time, bacterial species found in a healthy subgingival environment deserve further study to be able to predict the identification of new microbial periodontal pathogens.

3. Teodorescu AC, Martu I, Teslaru S, Kappenberg-Nitescu DC, Goriuc A, Luchian I, Martu MA, Solomon SM, Martu S. Assessment of Salivary Levels of RANKL and OPG in Aggressive versus Chronic Periodontitis. *J of Immunology Res*, Hindawi, 2019, Article ID 6195258, 6 pg. <https://doi.org/10.1155/2019/6195258>
4. Martu, I.; Goriuc, A.; Martu, M.A.; Vata, I.; Baciuc, R.; Mocanu, R.; Surdu, A.E.; Popa, C.; Luchian, I. Identification of Bacteria Involved in Periodontal Disease Using Molecular Biology Techniques. *Rev. Chim.* 2017, 68, 2407–2412
5. Solomon SM, Chiscop I, Moisei M, Badescu AC, Jelihovschi I, Martu-Stefanache A, Teusan A, Martu S, Iancu LS. Evaluation of DNA extraction. methods from saliva as a source of PCR - amplifiable genomic DNA. *Rev. Chim.*, 2015,67 (12):2101-2103
6. Martu, MA.; Maftai, GA.; Luchian, I.; Popa, C.; Filioreanu, AM.; Tatarciuc, D.; Nichitean, G.; Hurjui, LL.; Foia, LG. Wound healing of periodontal and oral tissues: Part II-Patho-physiological conditions and metabolic diseases. *Rom. J. Oral Rehabil.* 2020, 12, 30–40
7. Luchian I, Goriuc A, Sandu D, Covasa M. The Role of Matrix Metalloproteinases (MMP-8, MMP-9, MMP-13) in Periodontal and Peri-Implant Pathological Processes. *Int J Mol Sci.* 2022 Feb 4;23(3):1806.
8. Becker A, Zogakis I, Luchian I, Chaushu S. Surgical exposure of impacted canines: Open or closed surgery?. *Semin Orthod*, 2016; 22(1):27-33
9. Hiranmayi KV, Sirisha K, Ramoji Rao MV, Sudhakar P. Novel Pathogens in Periodontal Microbiology. *J Pharm Bioallied Sci.* 2017 Jul-Sep; 9(3):155-163. doi: 10.4103/jpbs.JPBS_288_16.
10. Mavrommatis K, Pukall R, (...) Kyrpides NC. Complete genome sequence of *Cryptobacterium curtum* type strain (12-3). *Stand Genomic Sci.* 2009 Sep 24;1(2):93-100. doi: 10.4056/sigs.12260.
11. Robertson D, Smith AJ. The microbiology of the acute dental abscess. *J Med Microbiol.* 2009 Feb;58(Pt 2):155-162. doi: 10.1099/jmm.0.003517-0. PMID: 19141730.
12. Uematsu H, Sato N, Djais A, Hoshino E. Degradation of arginine by *Slackia exigua* ATCC 700122 and *Cryptobacterium curtum* ATCC 700683. *Oral Microbiol Immunol.* 2006 Dec;21(6):381-4. doi: 10.1111/j.1399-302X.2006.00307.x.
13. Kumar PS, Griffen AL, Barton JA, Paster BJ, Moeschberger ML, Leys EJ. New bacterial species associated with chronic periodontitis. *J Dent Res.* 2003 May;82(5):338-44. doi: 10.1177/154405910308200503.
14. Nakazawa F, Poco SE, Ikeda T, Sato M, Kalfas S, Sundqvist G, Hoshino E. *Cryptobacterium curtum* gen. nov., sp. nov., a new genus of gram-positive anaerobic rod isolated from human oral cavities. *Int J Syst Bacteriol.* 1999 Jul;49 Pt 3:1193-200. doi: 10.1099/00207713-49-3-1193. Erratum in: *Int J Syst Bacteriol* 1999 Oct;49 Pt 4:1952.
15. Takeuchi Y, Umeda M, Sakamoto M, Benno Y, Huang Y, Ishikawa I (2001). *Treponema socranskii*, *Treponema denticola*, and *Porphyromonas gingivalis* are associated with severity of periodontal tissue destruction. *J Periodontol* 72:1354-1363.
16. Ximenez-Fyvie LA, Haffajee AD, Socransky SS (2000b). Microbial composition of supra- and subgingival plaque in subjects with adult periodontitis. *J Clin Periodontol* 27:722-732.
17. Popa, C.G.; Luchian, I.; Ioanid, N.; Goriuc, A.; Martu, I.; Bosinceanu, D.; Martu, M.A.; Tirca, T.; Martu, S. ELISA Evaluation of RANKL Levels in Gingival Fluid in Patients with Periodontitis and Occlusal Trauma. *Rev. Chim.* 2018, 69:1578–1580
18. Mocanu, RC.; Martu, M.A.; Luchian, I.; Sufaru, I.G.; Maftai, G.A.; Ioanid, N.; Martu, S.; Tatarciuc, M. Microbiologic Profiles of Patients with Dental Prosthetic Treatment and Periodontitis before and after Photoactivation Therapy—Randomized Clinical Trial. *Microorganisms* 2021, 9, 713.

19. Downes J, Munson MA, Spratt DA, Kononen E, Tarkka E, Jousimies-Somer H, Wade WG. Characterisation of Eubacterium-like strains isolated from oral infections. *J Med Microbiol.* 2001 Nov;50(11):947-951. doi: 10.1099/0022-1317-50-11-947.
20. Solomon, S.M.; Timpu, D.; Forna, D.A.; Stefanache, M.A.; Martu, S.; Stoleriu, S. AFM comparative study of root surface morphology after three methods of scaling. *Mater. Plast.* 2016, 53, 546–549
21. Solomon, S.M.; Stoleriu, S.; Forna, D.A.; Tampu, D.; Stefanache, M.A.; Ursarescu, I.G.; Martu, S. The Quantitative and Qualitative Assessment of Dental Substance Loss as Consequence of Root Planing by Three Different Techniques. *Mater Plast* 2016, 53:304-7.
22. Martu, M.A.; Maftei, G.A.; Luchian, I.; Stefanescu, O.M.; Scutariu, M.M.; Solomon, S.M. The Effect of Acknowledged and Novel Anti-Rheumatic Therapies on Periodontal Tissues—A Narrative Review. *Pharmaceuticals* 2021, 14, 1209. doi:10.3390/ph14121209
23. Ursu RG, Luchian I, Damian C, Porumb-Andrese E, Ghetu N, Cobzaru RG, Lunca C, Ripa C, Costin D, Jelihovschi I, Petrariu FD, Iancu LS. Diagnostics of HNSCC Patients: An Analysis of Cell Lines and Patient-Derived Xenograft Models for Personalized Therapeutical Medicine. *Diagnostics (Basel).* 2022 Apr 25;12(5):1071. doi: 10.3390/diagnostics12051071. PMID: 35626227; PMCID: PMC9139588.
24. Luchian, I.; Goriuc, A.; Martu, M.A.; Covasa, M. Clindamycin as an Alternative Option in Optimizing Periodontal Therapy. *Antibiotics* 2021, 10, 814. doi:10.3390/1007081425.
25. Laza GM, Sufaru IG, Martu MA, Martu C, Diaconu-Popa DA, Jelihovschi I, Martu S. Effects of Locally Delivered Minocycline Microspheres in Postmenopausal Female Patients with Periodontitis: A Clinical and Microbiological Study. *Diagnostics* 2022, 12, 1310. <https://www.mdpi.com/journal/diagnostics>
26. Spiridon I, Darie Nita RN, Kozlowski M, Nechita A, Ursu RG.: Influence of accelerated weathering on the performance of polylactic acid based materials, *Cellulose Chem Technol*, 2016, 50(5-6): 629-635,
27. Ursu RG, Iancu LS, Porumb-Andrese E, Damian C, Cobzaru RG, Nichitean G, Ripa C, Sandu D, Luchian I. Host mRNA analysis of periodontal disease patients positive for Porphyromonas gingivalis, Aggregatibacter actinomycetemcomitans and Tannerella forsythia, *Int J Mol Sci*, 2022, 23(17): art. no 9915.
28. Ursu RG, Luchian I, Ghetu, N, Costan VV, Stamatina O, Palade OD, Damian C, Iancu LS, Porumb-Andrese E. Emerging oncogenic viruses in head and neck cancers from romanian patients, *Appl Sci-Basel*, 2021, 11(19): art. no 9356.
29. Ursu RG, Giusca SE, Spiridon IA, Manole B, Danciu M, Costan VV, Palade DO, Ghetu N, Toader P, Vlad MA, Damian C, Porumb-Andrese E, Luchian I, Iancu LS. Understanding the Pattern of Oropharyngeal Cancers from North-East Romanian Patients. *Applied Sciences.* 2021; 11(24):12079. doi.org:10.3390/112412079
30. Ursu RG, Damian C, Porumb-Andrese E, Ghetu N, Cobzaru RG, Lunca C, Ripa C, Costin D, Jelihovschi I, Petrariu FD, Iancu LS. Merkel Cell Polyoma Virus and Cutaneous Human Papillomavirus Types in Skin Cancers: Optimal Detection Assays, Pathogenic Mechanisms, and Therapeutic Vaccination. *Pathogens.* 2022 Apr 16;11(4):479. doi: 10.3390/pathogens11040479. PMID: 35456154;
31. Nemescu D, Ursu RG, Nemescu ER, Negura L. Heterogeneous Distribution of Fetal Microchimerism in Local Breast Cancer Environment. *PLoS One.* 2016 Jan 25;11(1):e0147675. doi: 10.1371/journal.pone.0147675. PMID: 26808509;
32. Kumar C, et al.. Investigations into the polymorphisms at the ECM38 locus of two widely used *Saccharomyces cerevisiae* S288C strains, YPH499 and Y4742.

Yeast (2003)20(10):857-63

33. Lopez-Oliva I, Paropkari AD, Saraswat S, Serban S, Yonel Z, Sharma P, de Pablo P, Raza K, Filer A, Chapple I, Dietrich T, Grant MM, Kumar PS. Dysbiotic Subgingival Microbial Communities in Periodontally Healthy Patients With Rheumatoid Arthritis. *Arthritis Rheumatol.* 2018 Jul;70(7):1008-1013. doi: 10.1002/art.40485.
34. Anton DM, Martu A, Maris M, Maftei GA, Sufaru IG, Tatarciuc D, Luchian I, Ioanid N, Martu S. Study on the Effects of Melatonin on Glycemic Control and Periodontal Parameters in Patients with Type II Diabetes Mellitus and Periodontal Disease. *Medicina – Lithuania*, 2021, 57, 140. <https://doi.org/10.3390/medicina57020140>
35. Veisa G, Donciu M, Segal L, Hurjui L, Nistor I, Ursarescu I, Martu S, Burlea L, Solomon S. Albumin as a prognostic factor for malnutrition and inflammation in chronic kidney disease. *Rev. Chim. (Bucharest)*, 2016,67(1):103-105